



Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Ocular Oncology and Oculoplasty

Journal homepage: <https://ijooo.org/>



Guest Editorial

Pediatric ocular trauma

Mehul Shah^{1,*}

¹Founder and Director Drashti Netralaya, Dahod, Gujarat, India



ARTICLE INFO

Article history:

Received 20-10-2022

Accepted 27-10-2022

Available online 08-11-2022

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

Ocular trauma is an important topic with respect to preventable monocular blindness of the globe. Pediatric ocular trauma is a matter of great concern as sight loss in the younger age group puts extra economic and social burden as child has to live more blind years. Unresolved controversies and debatable management strategies need to be addressed with the development of novel research technology.¹

During current time of COVID lockdown schools are off rural children are more prone for injuries during outdoor play.²

Prospective, controlled, clinical studies cannot be conducted in ocular injury setting.

Children account for approximately one-third of the cases of severe eye injuries.^{2,3} However, the classification and scoring system in pediatric trauma is based on that developed for adults. The controversy over the position of zones 2 and 3 is pronounced in pediatric trauma.

Although management of trauma in children has many similarities to adults, striking differences are also noted. The adults achieve visual maturation, whereas amblyopia is a major contributor to poor outcomes in children, especially in those < 5-years-old.

When treating children with OGI, achieving a clear visual axis following globe repair must be accompanied

by accurate refraction and aggressive amblyopia therapy to improve the visual outcomes.⁴ Another difference is that a simple examination, especially in the early stages after trauma, can be difficult in a child requiring general anesthesia or sedation for proper assessment. Thus, the involvement of a pediatric ophthalmologist and access to pediatric facilities are essential for the management of OGI.

A major challenge in the care of eye trauma in children is the precise visual prognosis.

The pediatric ocular trauma score (POTS) down- played initial VA and added age and wound location as two new variables and excluded the afferent pupillary defect as a prognostic factor.⁵ Recently, the toddler/infant ocular trauma score (TOTS) was developed specifically for children < 6-years-old with traumatic OGIs.

For traumatic cataract we need to understand a few parameters, such as the age of the patient, expertise of the surgeon and assisting staff, availability of infrastructure facility, the status of the cataractous lens, and lens vitreous admixture that will guide the surgeon in planning the surgery, before effectuating primary or delayed cataract extraction and intraocular lens implantation. The increasing duration between injury and presentation may alter morphology because of absorption of lens matter and synechiae formation, which require different techniques and vitreoretinal interventions.⁶

* Corresponding author.

E-mail address: omtrustdahod@gmail.com (M. Shah).

A few studies compared visual outcome among open and closed globe injuries visual outcome, and was better in closed globe injuries.⁵ Whether single or multiple step surgeries are preferred is yet controversial in multistep management.⁷

One of the manifestations of ocular trauma is posttraumatic strabismus. The most common cause of alteration in motility is caused by the orbital fracture.^{8,9}

Pediatric Ocular Trauma is a complicated topic. Although several studies have helped resolve the underlying mysteries.

However, many multicenter studies are required to upgrade the knowledge and reduce unnecessary burden on society. An integrated trauma unit to coordinate various departments might be vital for the appropriate management of ophthalmic trauma in children.^{10,11}

Conflict of Interest

None.

References

1. Abrams GW, Kington RW. Falsely extinguished brightlight flash electroretinogram. Its association with dense vitreous hemorrhage. *Arch Ophthalmol.* 1984;100(9):1427–9. doi:10.1001/archophth.1982.01030040405005.
2. Shah M, Shah S, Patel H, Khanna R. Impact of Covid-19 lock down period on ocular trauma in pediatric population. *Int J Paediatr Geriatr.* 2020;3(2):27–29. doi:10.33545/26643685.2020.v3.i2a.86.
3. Acar U. A new ocular trauma score in pediatric penetrating eye injuries. *Eye.* 2011;25(3):370–4. doi:10.1038/eye.2010.211.
4. Adeoye A, Onakpoya OH. Indication for eye removal in Ile-Ife, Nigeria. *Afr J Med Med Sci.* 2007;36(4):371–5.
5. Shah M, Agrawal R, Teoh R, Shah SM, Patel K, Gupta S, et al. Pediatric ocular trauma score as a prognostic tool in the management of pediatric traumatic cataracts". *Graefe's Archive for. Graefes Arch Clin Exp Ophthalmol.* 2017;255(5):1027–36. doi:10.1007/s00417-017-3616-y.
6. Kuhn F. Traumatic cataract: what, when, how. *Graefes Arch Clin Exp Ophthalmol.* 2010;248(9):1221–3. doi:10.1007/s00417-010-1387-9.
7. Shah MA, Shah SM, Gosai SR, Gupta SS, Khanna RR, Patel KB, et al. Comparative study of visual outcome between open and closed-globe injuries following surgical treatment of traumatic cataract in children". *Eur J Ophthalmol.* 2018;28(4):406–11. doi:10.1177/1120672117747021.
8. Shah S, Shah M, Upadhyay P, Bardoloi G, Netralaya D. Strabismus in Cases of Cataract in Pediatric. *Open J Ophthalmol.* 2013;3(1):19–23. doi:10.4236/ojoph.2013.31006..
9. Roul-Yvonnet F, Tabchouri N. "[Children orbital floor fracture: retrospective study. *Ann Chir Plast Esthet.* 2012;57(3):240–4. doi:10.1016/j.anplas.2012.03.005.
10. Toh ZH, Agrawal S, Raje D, Hoskin A, Agrawal R, Khandelwal R, et al. International globe and adnexal trauma epidemiology study (IGATES): a report from Central India on visual outcome in open globe injuries and correlation with ocular trauma score. *Int Ophthalmol.* 2020;40(11):2797–806. doi:10.1007/s10792-020-01429-x.
11. Shah M, Bhasin P, Shah S, Patel H. Impact of Covid-19 lock down period on ocular trauma in pediatric population. *Int J Paediatr Geriatr.* 2020;3(2):27–9. doi:10.21203/rs.3.rs-49393/v1.

Author biography



Mehul Shah, Vitreo Retinal Surgeon
MS-FRF, FMRF
Founder and Director Drashti Netralaya, Dahod,
Gujarat, India
Email: omtrustdahod@gmail.com

Cite this article: Shah M. Pediatric ocular trauma. *IP Int J Ocul Oncol Oculoplasty* 2022;8(3):172-173.