

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Anatomy and Surgery of Head, Neck and Brain

Journal homepage: <https://www.ijashnb.org/>

Editorial

Epidemic keratoconjunctivitis: An overview

Rajendra Prakash Maurya^{1,*}, Aalok Kumar¹, Kokil Tiwari¹

¹Regional Institute of Ophthalmology, IMS, Banaras Hindu University, Varanasi, Uttar Pradesh, India



ARTICLE INFO

Article history:

Received 25-07-2023

Accepted 05-08-2023

Available online 08-08-2023

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

Epidemic keratoconjunctivitis (EKC) is a serious public health issue worldwide. EKC is highly contagious ocular surface infection mainly caused by adenoviruses. About 1% of all ophthalmic consultation in USA are related to conjunctivitis.¹ Adenovirus represents 15% to 70% of all infectious conjunctivitis.²

1. About the Virus

Human Adenovirus (HADVs) belongs to the family Adenoviridae, genus Mastadenovirus. It consists of around 130 serotypes out of which 54 types are pathogenic in humans. They are classified into 7 groups and are labeled from A to G. Adenoviruses are non-enveloped double-stranded DNA viruses. It's icosahedral capsids bears several specific antigens. HADV-8, HADV-19 & HADV-37 are responsible for contagious EKC.³

Adenoviruses are very stable and are highly resistant to chemical or physical agents even in adverse pH conditions. This leads to their prolonged survival outside the body and leading to increased chances of transmission. The contagious nature is best explained by considering data of HAdV19 which is viable for up-to 8 days on paper, 9 days on tonometer tips, 10 days on textiles and metal and up-to 35 days on plastics. Therefore, nosocomial spread is quite common in these infections.

* Corresponding author.

E-mail address: mauryarp_bhu@yahoo.com (R. P. Maurya).

2. Mode of transmission

Main transmitters of viruses are hands and contaminated objects like handkerchiefs, door-knobs, eye drops and medical instruments like tonometer & slit-lamp etc. Virus is transmitted through respiratory droplets and infected body fluids like tear-fluid which can enter the human body via nose, throat and conjunctiva. The viral incubation period ranges from 2-12 days.



Fig. 1: A female with bilateral EKC infection. Palpebral conjunctival hyperemia with follicles with eyelid edema.

3. Clinical presentations

EKC is an ocular surface infection which manifests as an intense inflammatory reaction with predominant clinical symptoms like redness, irritation / foreign body sensation, watery discharge, blurry vision and



Fig. 2: A boy with unilateral EKC infection with conjunctival edema and congestion with follicles with eyelid edema.

intolerance to light. The clinical signs are eyelid edema, follicular conjunctivitis, conjunctival hyperemia/congestion (Figures 1 and 2), sub-conjunctival hemorrhage (Figure 3), pseudomembrane formation (Figure 4) and often preauricular lymphadenopathy. The corneal involvement is also not uncommon. It can arise from the fourth day onward in the form of multifocal subepithelial infiltrates seen in the chronic stages of EKC. These small epithelial infiltrates tend to enlarge and develop into large nummuli that can lead to visual impairment. EKC presents as unilateral condition but in almost 70% of the cases it becomes bilateral in within a week of the presentation due to hand-to-eye transmission. It is a self limiting infection lasting for about 2 to 3 weeks.



Fig. 3: A young female having unilateral EKC infection with sub-conjunctival hemorrhage.

3.1. Diagnostic evaluations

It is mainly a clinical diagnosis, based on history and presenting signs and symptoms. The gold standard method for EKC diagnosis is detection of adenoviral DNA in conjunctival specimen by polymerase chain reaction (PCR) as it has been proven to be a very specific, sensitive and rapid test.⁴ The detection of adenovirus antigen by commercially available rapid tests are less sensitive than PCR technique.



Fig. 4: A man with unilateral EKC infection with membranous conjunctivitis having reduced vision.

3.2. Treatment

There is no definitive treatment for EKC, the treatment remains purely symptomatic as EKC is a self limiting disease. Cool compresses and artificial tears, supplemented with topical antibiotics to prevent or treat superinfection are the mainstay of treatment. The use of calcineurin inhibitors like topical cyclosporine-A has shows better results in treatment of chronic sub-epithelial infiltrates or nummuli.⁵ Some physicians prefer adding topical corticosteroids especially in cases of acute EKC where there is threat of incapacitating visual loss from persistent subepithelial infiltrates and its progression to pseudomembranous conjunctivitis and iridocyclitis.⁶ But steroids are double edged sword as even though it reduces inflammation and decreases corneal involvement but it has been found that it increases viral replication and titres.

There is no definitive antiviral treatment for EKC but certain drugs have reported clinical success like topical alpha-interferon, cidofovir, PVP-iodine 5%.⁷⁻⁹ Sundmacher et al found that Interferon eye drops might prevent infection in exposed persons.¹⁰

Off label use of antiseptic povidone-iodine ophthalmic solution is being employed as a means to decrease spread of EKC. It can rapidly inactivate adenovirus within 1 minute of exposure. Some studies have found that 2% solution can reduce the duration of infectivity.

4. Prevention

Basic control measures can be promoted from both patient's and hospital's side to prevent snowballing of the disease into an outbreak. Ophthalmic health care workers should always wear gloves while handling the patient. Proper hand hygiene should be maintained between examination of two patients along with frequent disinfection of instruments and surfaces. Employ mint of single-use disposable devices for examination if possible. Disinfection of the instruments with the help of isopropyl alcohol, 30% alcohol solution or sodium hypochlorite solution. Patient identification and isolation is a must to contain the spread. Health professionals should wear protective eyewear and should be asked to take leave if infected.¹¹

5. Conclusion

EKC is a very common ocular infection which should be dealt very cautiously with accurate diagnosis and prevention of transmission. A judicious case-by-case approach should be preferred to prevent drastic visual consequences.

6. Conflict of Interest

None.

7. Source of Funding

None.

References

- Smith AF, Waycaster C. Estimate of the direct and indirect annual cost of bacterial conjunctivitis in the United States. *BMC Ophthalmol.* 2009;9(1):13. doi:10.1186/1471-2415-9-13.
- Sohrabi M, Goodarzi Z, Saberfar E, Lashini H. The prevalence of viral conjunctivitis in patients who referred to eye specialist hospitals in Tehran, Iran. *Iran J Ophthalmol.* 2014;26(1):29–32.
- Erdin BN, Pas SD, Durak I, Schutten M, Saymer AA. A-5-year study of adenoviruses causing conjunctivitis in Izmir, Turkey. *J Med Virol.* 2015;87(3):472–7. doi:10.1002/jmv.24071.
- Sambursky R, Tauber S, Schirra F. The RPS adeno detector for diagnosing adenoviral conjunctivitis. *Ophthalmology.* 2006;113(10):1758–64. doi:10.1016/j.ophtha.2006.06.029.
- Reinhard T, Godehardt E, Pfahl HG, Sundmacher R. Local cyclosporine-A in nummuli after keratoconjunctivitis epidemica. A pilot study. *Ophthalmologe.* 2000;97(11):764–8. doi:10.1007/s003470070025.
- Kaufman HE. Adenovirus advances: new diagnostic and therapeutic options. *Curr Opin Ophthalmol.* 2011;22(4):290–3. doi:10.1097/ICU.0b013e3283477cb5.
- Lenaerts L, Naesens L. Antiviral therapy for adenovirus infections. *Antiviral Res.* 2006;17(2-3):172–80. doi:10.1016/j.antiviral.2006.04.007.
- Abel R, Abel AD. Use of PVP iodine in the treatment of presumptive adenoviral conjunctivitis. *Ann Ophthalmol Glaucom.* 1998;30:341–43.
- Meyer-Rüsenberg B, Loderstädt U, Richard G, Kaulfers PM, Gesser C. Epidemic Keratoconjunctivitis the current situation and recommendations for prevention and treatment. *Dtsch Arztebl Int.* 2011;108(27):475–80. doi:10.3238/arztebl.2011.0475.
- Sundmacher R, Wigand R, Cantell K. The value of exogenous interferon in adenovirus keratoconjunctivitis. *Graefes Arch Clin Exp Ophthalmol.* 1982;218(3):139–40. doi:10.1007/BF02215650.
- Sambursky RP, Fram N, Cohen EJ. The prevalence of adenoviral conjunctivitis at the Wills Eye Hospital emergency room. *Optometry.* 2007;78(5):236–9. doi:10.1016/j.optm.2006.11.012.

Author biography



Rajendra Prakash Maurya, Associate Professor,
Institute of Medical Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh, India
<https://orcid.org/0000-0001-9343-6003>

Aalok Kumar, Assistant Professor

Kokil Tiwari, Junior Resident

Cite this article: Maurya RP, Kumar A, Tiwari K. Epidemic keratoconjunctivitis: An overview. *IP Indian J Anat Surg Head, Neck Brain* 2023;9(2):53-55.