

Management of orbital disorders is a challenging job

Rajendra P Maurya
Editor in Chief IJCEO
Assistant Professor & I/c Orbit, Ocular
Oncology and Oculoplasty Unit
Department of Ophthalmology,
Institute of Medical Sciences,
Banaras Hindu University, Varanasi, (UP), INDIA
E-mail: editorijceo@gmail.com, mauryarp_bhu@yahoo.com

Dear Friends,

Management of orbital disorders is a great challenge for any one. Orbit is the only cavity having small size with wide diversity of disorders. The presence of complex, multiple anatomical structures arisen from all three embryonic germ layers is responsible for highly diversified clinical presentation thus making differential diagnosis quite difficult. Orbit is very small in volume (30cc) and enclosed space (posteriorly by four bony walls and anteriorly by Orbital septum) with much less capacity to adjust anatomical and physiological alterations following orbital trauma and space occupying lesions resulting in comportment syndrome. Orbital comportment syndrome is a true ocular emergency need immediate identification and intervention to be avoided for permanent visual loss.⁽¹⁾

Orbit can be affected from disease from neighboring structures (nose, paranasal sinuses, pterygopalatine fossa and cranial cavity etc.) through multiple communications like fissures, canal and foramena. Being highly vascularized area, orbit may get metastasis from distant malignant neoplasm. Developmental disorders of orbit often occurs in combination with craniofacial malformations that usually required multiple staged larger surgeries.

Detailed history, extensive clinical examination and modern imaging tools (ultrasonography, multislice CT Scanner & MRI etc.) is helpful in narrowing differential diagnosis. (2) Newer pathological techniques like fine needle aspiration biopsy & special immunohistochemistry also helpful for accurate diagnosis. Thus pathologist is a valuable resource for Orbitologist but it is necessary to communicate with the pathologist before and after surgical procedures to provide accurate patient care. Orbital navigation technique is another boon for orbitologist in managing orbital disorders. Management of orbital disorders requires knowledge of comprehensive anatomy of orbit, additional expertise and dedicated team of other specialist like neurosurgeon, otolaryngologists and maxillofacial surgeons.

References

- 1. Linberg JV. Orbital compartment syndromes following trauma. Adv Ophthalmic Plast Reconstr Surg 1987;6:51-62.
- Zuravleff JJ Johnson MH. An ophthalmic surgeons view of orbital imaging techniques. Semin Ultrasound CT, MR 1997;18(6):395-402.
- 3. Agrawal P, Dey P. Lal A. Fine -needle aspiration cytology of orbital and eyelid lesions. Diag Cytopathol 2013;41(11):1000-11.