



## Original Research Article

## Changes in keratometry and refractive status pre and post pterygium surgery

Bansari Vadodaria<sup>1</sup>, Aishwariya Thakre<sup>1</sup>, Rupali Maheshgauri<sup>1,\*</sup>, Divya Motwani<sup>1</sup>,  
Ashish Mishra<sup>1</sup>

<sup>1</sup>Dept. of Ophthalmology, Dr. D. Y. Patil Medical College, Hospital & Research Centre, Pimpri, Pune, Maharashtra, India



## ARTICLE INFO

## Article history:

Received 18-11-2019

Accepted 15-12-2019

Available online 26-12-2019

## Keywords:

Keratometry

Changes in keratometry readings in pterygium

Pterygium induced refractive error

Pterygium induced corneal astigmatism

Sutureless pterygium surgery

Post pterygium refractive error

## ABSTRACT

**Purpose:** To study pterygium induced astigmatism. In this study, we have analyzed our results of refractive status of pterygium patient after pterygium excision surgery. Up till now we have seen pterygium induced astigmatism but with the help of this study we would like to find out refractive status of patient post operatively which will help in refractive cataract surgery of patient having both pterygium and cataract.

**Materials and Methods:** Descriptive study was undertaken of 50 cases of patients diagnosed to have pterygium induced astigmatism. Patients were explained about the study after which written and informed consent was taken about their participation in the study. Demographic factors like age, sex, occupation and address were recorded. Complete ophthalmic and medical history was taken, Visual acuity using Snellen's chart, Refractive status was deduced. Keratometry observed for horizontal and vertical diameters corneal diopter by bausch and lomb keratometry. Pterygium induced astigmatism which were operated by pterygium excision and conjunctival autograft with suture or without suture. Patients were examined on post operative day 1,7,21 for refractive status where the refractive status of patients with pterygium pre-operative and post operative with conjunctival autograft with suture and without suture was evaluated. Data was entered in Microsoft Excel and statistical analysis was done using software EpiInfo or SPSS. Quantitative data was analysed in terms of Means and Standard Deviation. Qualitative data was analysed with appropriate test of significance like Unpaired T test and Paired T test. The probability level of 0.05 was considered as statistically significant.

**Result:** There was significant difference was seen in mean KV and KH pre-operatively, on post operative day 1 after pterygium excision surgery with conjunctival autograft with or without suture and on regular follow up ( $p = 0.000$ ), such that KV decreases while KH increases following surgery and serial follow up. Mean (KV minus KH) pre-operatively, on post operative day 1 and on regular follow up was statistically significant ( $p = 0.000$ ). (KV minus KH) decreases after pterygium excision surgery with conjunctival autograft with or without suture. Spherical and Astigmatic error reduces significantly after pterygium excision surgery with conjunctival autograft with or without suture ( $p = 0.000$ ).

**Conclusion:** Pterygium leads to significant high corneal astigmatism, which hampers vision of the patient. Pterygium causes horizontal meridian flattening (KH). We concluded that, pterygium causes simple myopic with the rule type of astigmatism and compound hypermetropic and myopic astigmatism type of refractive error. With the rule astigmatism was most common pre-operatively. There is drastic changes in keratometry readings pre-operatively and post surgical excision of pterygium tissue. Overall, after pterygium excision surgery most of the patients had no refractive error except simple myopic type of refractive error. After pterygium excision surgery astigmatism was reduced, in compound astigmatic type of refractive error. There is definite change in refractive status in pterygium post pterygium surgery so with this study I would like to deduce that, to get emmetropic status for cataract surgery or refractive cataract surgery pterygium tissue excision should be done.

© 2019 Published by Innovative Publication. This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by/4.0/>)

## 1. Introduction

Pterygium is a triangular fibrovascular subepithelial ingrowth of degenerative bulbar conjunctival tissue over the limbus onto the cornea.<sup>1</sup> Pterygium mostly causes "with the rule astigmatism" that is: vertical meridian is more steeper than horizontal meridian as pterygium generally encroaches the cornea from the nasal side in a horizontal direction at first which causes the H meridian to flatten but gradually when the pterygium progresses it leads to irregular astigmatism.<sup>2</sup>

Grading of pterygium according to Youngson R.M<sup>3</sup>  
Grades of pterygium

1. Pterygium invading < 1.5 mm of cornea
2. Pterygium invading < half the radius of cornea
3. Pterygium invading > half the radius of cornea
4. Pterygium almost or reaching the centre of cornea

## 2. Aims and Objectives

### 2.1. Aim

To study pterygium induced astigmatism and post operative refractive status of pterygium patient.

### 2.2. Objectives

1. To evaluate pterygium induced astigmatism.
2. Analysis of the pterygium size inducing marked refractive astigmatism.
3. To evaluate post operative refractive status.

## 3. Review of Literature

Three topographic properties of the cornea are important to its optical function: the underlying shape, which determines its curvature and hence its refractive power.<sup>4</sup>

The development of a pterygium can lead to significant astigmatism. A pterygium generally causes localized flattening central to the apex of the pterygium.<sup>5</sup> As this flattening is along the horizontal meridian, it usually causes with-the-rule corneal astigmatism.<sup>6</sup> The vertical corneal meridian is steep in younger adults. This reduces with age and tends to give rise to against the rule astigmatism in later years.<sup>7</sup> Fong et al. in 1998 observed that pterygium excision usually induces a reversal of pterygium-related corneal flattening.<sup>8</sup>

Pterygium is described as an active, invasive and inflammatory process where subconjunctival tissue undergoes elastotic degeneration and proliferates as fibrovascular granulation tissue under the epithelium, which encroaches onto cornea.<sup>9</sup> Pterygium is an acquired, chronic and degenerative disease. The exact etiology of pterygium is still unknown. But, it is more common in people living in

hot dry climates. Due to prolonged effect of environmental factors such as exposure to sunlight, dry heat, high wind and abundance of dust. It is more common in males than female because of long outdoor work. Pterygium usually occurs on the nasal side of bulbar conjunctiva between inter palpebral fissures. It can also occur at temporal side of bulbar conjunctiva but it is more commonly seen along with nasal pterygium. Both Nasal and temporal pterygium in one eye is called as Double Pterygium. Pterygium can be unilateral or bilateral, but asymmetrical. It is common in old age. Pterygium presents mainly with symptoms like foreign body sensation, irritation, cosmetic reason, impairment of vision as it encroaches onto cornea and causes astigmatism. Pterygium can progress and can reach up to pupillary margin or beyond that covering pupil and causes visual axis obstruction of visual axis.<sup>10</sup>


Flattening was seen in the horizontal meridian, which was associated with astigmatism. The exact mechanism of flattening is not clear. It is thought to be caused by the formation of tear meniscus between the corneal apex and the elevated pterygium, causing an apparent flattening of the normal corneal curvature. Pterygium induces significant astigmatism which may be either 'with-the-rule' (WTR) or 'against-the-rule' (ATR). The mechanisms explaining the astigmatism are: the tractional force of contractile elements within the pterygium lead to mechanical distortion and flattening of the cornea in its horizontal meridian leading to hypermetropic WTR astigmatism. Pterygium induced with-the-rule corneal astigmatism is hemimeridional on the side of the pterygium resulting in a localized flattening of the cornea central to the leading apex. The localised pooling of tears at the advancing edgehead of the pterygium is also responsible for corneal flattening. Pterygium results in high corneal astigmatism, which decreases following an excision. Pterygium can be divided into 4 parts:- 1) Head: - Apical part of pterygium on the cornea. 2) Neck: - lies between head and body onto the limbus. 3) Body: - It represents as main part of the pterygium lying over the sclera and extending from the canthal side of bulbar conjunctiva. 4) Cap: - It is present just in front of the head. It can be of two types, few whitish infiltrates seen just in front of head of pterygium is called "Fuch's spots", "Fuch's islands" or "Fuch's islets" and another is "Stocker's line" - an iron deposited line in front of head of pterygium.

According to the progression pterygium is of two types:- (1) PROGRESSIVE: - It is described as thick, fleshy and vascular pterygium with fuch's spot. (2) REGRESSIVE OR ATROPHIC: - thin, atrophic, attenuated pterygium with very little vascularity. Stocker's line is present in this type of pterygium. This pterygium ultimately regresses and becomes more atrophic and membranous but it never disappears.<sup>11</sup>

Pterygium more frequently occur on nasal side than temporal side. The bulbar conjunctiva of temporal side

\* Corresponding author.

E-mail address: [rupalim021@gmail.com](mailto:rupalim021@gmail.com) (R. Maheshgauri).




**Fig. 1:** Parts of pterygium


of the cornea is situated below the lacrimal gland so it is less prone to drying as it is bathed by fresh tears so the relative rarity of pterygium is in the temporal side. Rarity of temporal side of pterygium can be explained by aqueous tear secretion but it does not explain mucin deficiency association.<sup>11</sup> Pterygium present on the nasal side is an indication of the pathogenetic role of the solar light as light gets reflected from lateral wall of nose onto the nasal side of bulbar conjunctiva. Nasal pterygium can also be explained by noting that longer temporal eyelash of upper eyelid shade and filter sunlight falling on the temporal conjunctiva and cornea than on nasal side.<sup>12,13</sup> Elliot explained that because of wind, drying tissue of the medial third of the palpebral aperture devitalizes which leads to actinic reaction to damage cornea and conjunctival layers.<sup>14</sup> pterygium occurs more in dominant eye, probably due to the non-dominant eye is closed in intense sunlight.<sup>13</sup>



**Fig. 3:** Nasal atrophic pterygium



**Fig. 2:** Nasal progressive pterygium



**Fig. 4:** Double head pterygium

Different surgical options for pterygium are:

1. Bare sclera excision (d'ombrain)
2. Mitomycin (mmc)
3. Pterygium excision with amniotic membrane transplant
4. Pterygium excision with conjunctivo-limbal autograft transplantation using fibrin glue
5. Pterygium excision surgery with conjunctivo -limbal autograft transplantation using sutures
6. Pterygium excision with conjunctivo-limbal autograft transplantation using patient's own blood

7. Lamellar keratoplasty and excimer laser phototherapeutic keratectomy (are being studied)

Out of all techniques conjunctival autografting is the best available option after pterygium excision to prevent recurrence.<sup>15–18</sup> Study has compared sutureless glue-free versus sutured limbal conjunctival autograft in primary pterygium surgery in adults and showed sutureless glue-free limbal conjunctival autograft is safe, effective, economical and had greater patient satisfaction.<sup>18</sup>

#### 4. Materials and Methods

A comparative study was undertaken where the refractive status of patients with pterygium pre-operative and post operative with conjunctival autograft with suture and without suture was evaluated.

##### 4.1. Study area

D.Y. Patil Medical College, Pimpri, Pune.

##### 4.2. Study design

Descriptive study

##### 4.3. Study period

September to September-2019

Institute ethics committee clearance have been obtained before the start of the study.

Written and informed consent was obtained from all the patients.

##### 4.4. Sample size

50 cases

Considering the average number of patient attending the OPD in D.Y. Patil medical college and having diagnosed pterygium being 2-3 cases per month :

The period of the study being 24 month, sample size was 50 cases.

##### 4.5. Inclusion criteria

1. Pterygium patients age between: 20-40
2. Pterygium patients without history of spectacles.

##### 4.6. Exclusion criteria

1. Pterygium patients with previous history of spectacles. i.e refractive error.
2. Pterygium patients with history of recurrent pterygium.
3. Pseudopterygium.

#### 5. Methodology

All patients included in the study were diagnosed to have pterygium induced astigmatism. Patients were explained about the study after which written and informed consent was taken about their participation in the study.

Following protocols were undertaken in each case:

1. Demographic factors like age, sex, occupation and address were recorded as per the attached proforma.
2. Complete ophthalmic and medical history was taken
3. Visual acuity using Snellen's chart.
4. Refractive status deduced
5. Keratometry observed for horizontal and vertical diameters corneal dioptric

##### 5.1. Surgical technique.

Total of 50 cases of pterygium induced astigmatism were included in this study which were operated by pterygium excision and conjunctival autograft with suture or without suture.

Patients were examined on post operative day 1,7,21 for refractive status.

##### 5.2. Statistical analysis

Data was entered in Microsoft Excel and statistical analysis was done using software EpiInfo or SPSS.

Quantitative data was analysed in terms of Means and Standard Deviation.

Qualitative data was analysed with appropriate test of significance like Unpaired T test and Paired T test.

#### 6. Results

Total of 50 cases of pterygium induced astigmatism were included in this study. Out of which 25 patients were operated with conjunctival autograft without suture and 25 patients were operated using conjunctival autograft with suture technique. These were following observations and results of the present study:

**Table 1:** Size of pterygium (mm) Statistics

N	50.00
Mean	2.52
Std. Error of Mean	0.15
Std. Deviation	1.06
Range	3.50
Minimum	1.00
Maximum	4.50

In present study mean size of pterygium tissue was 2.52 mm with standard deviation of 1.06 mm, with the highest 4.5 mm and lowest 1 mm.

In present study Grade 3 (44%) was most common followed by grade 2 (26%) followed by grade 4 (24%).




Chart 1: Grade of pterygium




Chart 2: Co-relation of grade of pterygium with keratometry readings (KV-KH)

On application of spearman’s correlation, there was positive correlation between grade of pterygium and KV – KH (diopter) which was highly significant. It means as grade of pterygium increases, difference between KV – KH also increases (Spearman’s rho 0.482, p = 0.000)

Table 3 statistics shows that, significant difference was seen in mean KV and KH pre-operatively, on post operative day 1 after pterygium excision surgery with conjunctival autograft with or without suture and on regular follow up (p = 0.000), such that KV(43.84 + 1.01 decreases to 43.08 + 0.72) decreases while KH (41.30 + 0.96 increases to 42.56 + 0.79) increases following surgery and serial follow up.

Table 4 statistics shows that, mean (KV minus KH) pre-operatively, on post operative day 1 and on regular follow

up was statistically significant (p = 0.000). (KV minus KH) decreases (2.55 + 1.14 reduces to 0.60 + 0.41) after pterygium excision surgery with conjunctival autograft with or without suture

The present study shows that, mean spherical error (SE) (1.06 + 1.39) and astigmatic error (AE) (1.02 + 0.89) pre-operatively, on post operative day 1 and on regular follow up was statistically significant (p = 0.000). Spherical (0.08 + 0.23) and Astigmatic error (0.17 + 0.26) reduces significantly after pterygium excision surgery with conjunctival autograft with or without suture.

On application of unpaired t test, no significant difference was observed between mean vertical (KV) and horizontal (KH) or their difference (KV-KH) by two types of surgery (Pterygium excision with conjunctival autograft without suture or with suture) at various duration of study (p > 0.05). It mean in comparison between pre-operative and post-operative status there was equal reduction of KV, difference between KV & KH, increase in KH following two types of surgery.

Chart 3 shows that, simple myopic astigmatism (32%) was most common pre-operatively followed by compound hypermetropic astigmatism (28%). Post-operatively, 56% subjects were having no refractive error followed by simple myopic astigmatism (26%). Almost no change was seen in spherical refractive error where as astigmatic error was reduced or nullified.

Chart 4 shows that, with the rule astigmatism (58%) was most common pre-operatively followed by against the rule astigmatism (32%). Post-operatively, majority of the subjects that is (68%) subjects were having no astigmatism while only (14%) patients had against the rule astigmatism.

Chart 5 shows that, compound hypermetropic astigmatism was seen in most of the cases of grade 3 and 4 pterygium and simple hypermetropic astigmatism was seen in grade 2 pterygium. Grade 1 causes only simple myopic astigmatism.

Chart 6 shows that, after pteryguim excision surgery patients with grade 3 and grade 2 pterygium patients had drastic reduction in astigmatic error.

Comparing pre-operative and post-operative refractive error after pterygium surgery shows that, in grade 1 pterygium simple myopic astigmatic error almost remains same. In Grade 2, 3 pterygium patients had no refractive error to mild simple myopic astigmatic error but compound hypermetropic astigmatism is completely nullified.

## 7. Discussion

In present study Grade 3 (44%) was most common followed by grade 2 (26%) followed by grade 4 (24%). In present study mean length of pterygium tissue was 2.52 mm with standard deviation of 1.06 mm, with the highest 4.5 mm and lowest 1 mm. There was positive correlation between grade of pterygium and KV – KH (diopter) and it was

**Table 2:** Keratometry readings in vertical (KV) and horizontal (KH) meridian pre-operatively and post-operatively

Time	Group	Mean	N	Std. Deviation	Std. Error Mean
Pre-Operative	K V	43.84	50.00	1.01	0.14
	K H	41.30	50.00	0.96	0.14
1 Day	K V	43.37	50.00	0.88	0.12
	K H	42.09	50.00	0.78	0.11
1 Week	K V	43.06	50.00	0.76	0.11
	K H	42.29	50.00	0.83	0.12
3 Week	K V	43.08	50.00	0.72	0.10
	K H	42.56	50.00	0.79	0.11

**Table 3:** Comparison of KV and KH pre-operatively and post-operatively

Paired samples test								
Pair	Mean	Std. Deviation	Std. Error Difference	95% CI of the Difference		t	Df	Sig. (2-tailed)
				Lower	Upper			
KV – KV 1 Day	0.47	0.62	0.09	0.29	0.65	5.36	49.00	0.00
KV – KV 1 Week	0.79	0.59	0.08	0.62	0.95	9.39	49.00	0.00
KV – KV 3 Week	0.77	0.54	0.08	0.61	0.92	9.97	49.00	0.00
KH – KH 1 Day	-0.80	0.94	0.13	-1.06	-0.53	-5.99	49.00	0.00
KH – KH 1 Week	-1.00	0.98	0.14	-1.27	-0.72	-7.16	49.00	0.00
KH – KH 3 Week	-1.27	0.99	0.14	-1.55	-0.99	-9.07	49.00	0.00

**Table 4:** Mean (KV minus KH) pre-operatively, on post operative day 1 and on regular follow up

Difference between KV and KH (KV-KH) (diopter)								
Time	Group	Mean	N	Std. Deviation	Std. Error Mean			
Pre-Operative	KV – KH	2.55	50.00	1.14	0.16			
1 Day	KV – KH	1.28	50.00	0.84	0.12			
1 Week	KV – KH	0.89	50.00	0.65	0.09			
3 Week	KV – KH	0.60	50.00	0.41	0.06			
Paired samples test								
Pair	Mean	Std. Deviation	Std. Error Difference	95% CI of the Difference		T	Df	Sig. (2-tailed)
				Lower	Upper			
KV - KH 1 Day	1.27	1.09	0.15	0.95	1.58	8.18	49.00	0.00
KV – KH 1 Week	1.66	1.10	0.16	1.34	1.97	10.63	49.00	0.00
KV - KH 3 Week	1.94	1.11	0.16	1.62	2.26	12.35	49.00	0.00

highly significant. It means as grade of pterygium increases, difference between KV-KH also increases (Spearman's rho 0.482,  $p = 0.000$ ). As, difference between KV-KH increases astigmatism increases compound hypermetropic astigmatism was seen in most of the cases of grade 3 and 4 pterygium, simple hypermetropic astigmatism was seen in grade 2 pterygium and Grade 1 pterygium causes only simple myopic astigmatism. After pterygium surgery patients with grade 3 and grade 2 patients had drastic reduction in astigmatic error. Comparing pre-operative

and post-operative refractive error after pterygium excision surgery shows that, in grade 1 pterygium simple myopic astigmatic error almost remains same. In Grade 2, 3 pterygium patients had no refractive error to mild simple myopic astigmatic error but compound hypermetropic astigmatism is completely nullified so patient with grade > 2 pterygium should undergo pterygium excision surgery. Similar study was carried out by Poonam Bhargava<sup>19</sup> on 52 eyes of 50 patients with primary pterygium (32.69%) had grade III pterygium was seen in (32.69%), grade I (23.07%)

**Table 5:** Change in refractive error pre-operatively and on post-operative day 1, 1 week, 3 week

Refractive error (diopter)					
Time	Group	Mean	N	Std. Deviation	Std. Error Mean
Pre-Operative	SE	1.06	50.00	1.39	0.20
	AE	1.02	50.00	0.89	0.13
1 Day	SE	0.12	50.00	0.30	0.04
	AE	0.50	50.00	0.40	0.06
1 Week	SE	0.06	50.00	0.24	0.03
	AE	0.26	50.00	0.34	0.05
3 Week	SE	0.08	50.00	0.23	0.03
	AE	0.17	50.00	0.26	0.04

**Table 6:** Change in spherical and astigmatism error pre-operatively and post-operatively

Paired samples test								
Pair	Mean	Std. Deviation	Std. Error Difference	95% CI of the Difference		t	Df	Sig. (2-tailed)
				Lower	Upper			
SE - SE 1 Day	0.95	1.21	0.17	0.60	1.29	5.51	49.00	0.00
SE - SE 1 Week	1.00	1.32	0.19	0.63	1.38	5.36	49.00	0.00
SE - SE 3 Week	0.98	1.31	0.19	0.61	1.35	5.30	49.00	0.00
AE – AE 1 Day	0.53	0.73	0.10	0.32	0.73	5.12	49.00	0.00
AE – AE 1 Week	0.76	0.85	0.12	0.52	1.00	6.35	49.00	0.00
AE – AE 3 Week	0.85	0.79	0.11	0.63	1.07	7.62	49.00	0.00




Chart 3: Change different type of refractive errors among patients pre-operatively and post-operatively

**Table 7:** Comparison of keratometry readings between two type of surgery

Time	Group	Surgery	Mean	Std. Deviation	Std. Error Mean	P value
Pre-Operative	K V	1	43.76	0.99	0.20	0.58
		2	43.92	1.04	0.21	
	K H	1	41.30	0.85	0.17	0.97
		2	41.29	1.08	0.22	
	KV - KH	1	2.46	1.00	0.20	0.60
		2	2.63	1.28	0.26	
1 Day	K V	1	43.39	0.91	0.18	0.87
		2	43.35	0.86	0.17	
	K H	1	42.05	0.76	0.15	0.72
		2	42.13	0.82	0.16	
	KV - KH	1	1.34	0.88	0.18	0.62
		2	1.22	0.82	0.16	
1 Week	K V	1	42.94	0.70	0.14	0.29
		2	43.17	0.82	0.16	
	K H	1	42.26	0.81	0.16	0.80
		2	42.32	0.86	0.17	
	KV - KH	1	0.89	0.62	0.12	1.00
		2	0.89	0.70	0.14	
3 Week	K V	1	43.00	0.70	0.14	0.47
		2	43.15	0.74	0.15	
	K H	1	42.43	0.77	0.15	0.25
		2	42.69	0.81	0.16	
	KV - KH	1	0.69	0.41	0.08	0.15
		2	0.52	0.40	0.08	

Pterygium excision with conjunctival autograft without suture = 1  
Pterygium excision with conjunctival autograft with suture = 2




Chart 4: Change in type of astigmatism among patients pre-operatively and on regular follow up post-operatively






Chart 5: Co-relation between grade of pterygium and pre-operative refractive error




Chart 6: Co-relation between grade of pterygium and post-operative refractive error

and grade II (23.07%), grade IV was seen in 21.15%. Dr. Eknath Shelke<sup>20</sup> found that out of 37 patients operated for pterygium with conjunctival autograft, 33 eyes belonged to Grade II and III pterygium (89.19%) only 2.70% had Grade I and 8.11% had Grade IV pterygium seen, which has a significant impact on vision. Maheshwari also got same results where 36 eyes had primary pterygium, where Grade II type of pterygium was most common i.e., 44.45% and Grade III pterygium was 33.33%. Similar observations were found in the study done by Fong<sup>8</sup> and Avisar.<sup>21</sup> Thus as the grade increases there is increase in astigmatic refractive error and hence the patient seeks for remedy.

In present study mean size of pterygium tissue was 2.52 mm with standard deviation of 1.06 mm, with the highest 4.5 mm and lowest 1 mm. As the size of pterygium increases it

makes the horizontal meridian flatter causing KH to increase there by increasing difference between keratometry readings (KV-KH) leading to increase in astigmatic refractive error. Lesions extending >45% of the corneal radius or within 3.2 mm of the visual axis produce increasing degrees of induced astigmatism.<sup>22</sup> The authors reported that pterygium with larger than 2.2 mm extension might contribute to corneal astigmatism >2 D.<sup>23</sup> Avisar A et al<sup>21</sup> noted that as primary pterygium encroaches more than 1.00 mm size from limbus it causes with-the-rule significant astigmatism (> or = 1.0 diopter). Kampitak noted that pterygium induced corneal astigmatism and timing of pterygium excision surgery are associated with size of pterygium. He also noted that about 2 D of astigmatism is caused by 2.25mm of pterygium which should be considered as the limit of surgery.<sup>24</sup>

The present study shows that, mean spherical error (SE) ( $1.06 + 1.39$ ) and astigmatic error (AE) ( $1.02 + 0.89$ ) pre-operatively, on post operative day 1 and on regular follow up was statistically significant ( $p = 0.000$ ). Spherical ( $0.08 + 0.23$ ) and Astigmatic error ( $0.17 + 0.26$ ) reduces significantly after pterygium excision surgery with conjunctival autograft with or without suture. Similar results were found in study done by Popat KB et al<sup>23</sup> where Mean astigmatism preoperatively was found to be  $6.20 \pm 3.58$  D which subsequently decreased to  $1.20 \pm 1.27$  D on 45th post-operative day-showing  $5.09 \pm 3.32$  D of change in astigmatism which was statistically significant (paired t-test,  $p < 0.05$ ). In the study done by Maheshwari S.<sup>25</sup> corneal astigmatism reduced from  $4.40 \pm 3.64$  diopter (D) to  $1.55 \pm 1.63$  D (P value  $< 0.001$ ) following surgery, which is comparable to our study. Also Stern and Lin reported improvement in topographic indices in 16 eyes ; they reported corneal astigmatism to reduce from ( $5.93$ – $2.46$  D) to ( $1.92$ – $1.68$  D).

The present study shows that with the rule astigmatism (58%) was most common pre-operatively followed by against the rule astigmatism (32%). Post-operatively, 68% subjects were having no astigmatism followed by against the rule astigmatism (14%). Similar results were found in study done by Eias Ali RT, Mohamed Ali AB on 100 Sudanese males and females aged between 18 and 45 years having pterygium in one eye and another eye being normal that The primary pterygium caused meridional steepness in corneal curvature (98%). With-the-rule corneal astigmatism was dominant (61%) ( $p < 0.001$ ). This result reveals that primary pterygium often leads to steepness in vertical meridian rather than horizontal meridian. This condition occurs because pterygium generally invade the cornea from the nasal side in a horizontal direction and this causes the horizontal meridian to flatten.<sup>26</sup> Similar result was found by Avisar et al.<sup>24</sup> Study carried out by Stern and Lin also suggested that pterygia induce asymmetric with-the-rule astigmatism.<sup>22</sup>

Present study shows, significant difference was seen in mean KV and KH pre-operatively, on post-operative day 1 after pterygium excision surgery with conjunctival autograft with or without suture and on regular follow up ( $p = 0.000$ ), such that KV ( $43.84 + 1.01$  decreases to  $43.08 + 0.72$ ) decreases while KH ( $41.30 + 0.96$  increases to  $42.56 + 0.79$ ) increases following surgery and serial follow up. Similar results were seen by S. Maheshwari flattening was seen in the horizontal meridian, which was associated with astigmatism. It is thought to be caused by the formation of tear meniscus between the corneal apex and the elevated pterygium, causing an apparent flattening of the normal corneal curvature.<sup>27</sup>

The present study shows that, mean (KV minus KH) pre-operatively, on post operative day 1 and on regular follow up was statistically significant ( $p = 0.000$ ). (KV

minus KH) decreases ( $2.55 + 1.14$  reduces to  $0.60 + 0.41$ ) after pterygium excision surgery with conjunctival autograft with or without suture, leading to reduction in refractive error i.e mean spherical error (SE) ( $1.06 + 1.39$ ) and astigmatic error (AE) ( $1.02 + 0.89$ ) pre-operatively reduced to Spherical ( $0.08 + 0.23$ ) and Astigmatic error ( $0.17 + 0.26$ ) post-operatively day 21. ( $p = 0.000$ ) We found that Post-operatively, 56% subjects were having no refractive error followed by simple myopic astigmatism (26%). Post-operatively, 68% subjects were having no astigmatism followed by against the rule astigmatism (14%). Fong et al. in 1998 observed that pterygium excision usually induces a reversal of pterygium-related corneal flattening.<sup>8</sup>

In present study we found that simple myopic astigmatism (32%) was most common pre-operatively followed by compound hypermetropic astigmatism (28%). Post-operatively, 56% subjects were having no refractive error followed by simple myopic astigmatism 26%. With the rule astigmatism 58% was most common pre-operatively followed by against the rule astigmatism 32%. Post-operatively, 68% subjects were having no astigmatism followed by against the rule astigmatism 14%. In Grade 2, 3 pterygium patients had no refractive error to mild simple myopic astigmatic error but compound hypermetropic astigmatism is completely nullified. Other similar study done by Lin A, Stern GA. found that hypermetropic WTR astigmatism was but common.<sup>22</sup> Pterygium results in high corneal astigmatism, which decreases following an excision.<sup>28</sup> A pterygium-induced refractive change often leads to visual impairment, Keratometry measures only the central cornea and peripheral cornea is ignored and hence the results can be erroneous in eyes with pterygium.<sup>27</sup> So, Computerized videokeratography remains the best tool in evaluating pterygium-associated corneal changes.<sup>27</sup>

In present study, the patient compliance On application of Mann-Whitney U test, pain score (done on basis of Wong Baker faces pain rating scale) after pterygium excision with conjunctival autograft without suture was lower than pain score in pterygium excision with conjunctival autograft with suture and difference between the two surgeries was statistically highly significant ( $p = 0.00$ ).

There was no significant difference seen in post - operative after effects between two surgical techniques among patients ungergoing pterygium excision surgery. According to Dr. Mitra,<sup>29</sup> “The main disadvantage of sutureless glue-free technique is the risk of graft loss in the immediate postoperative period.” In present there was 1 case of graft lost in sutureless technique and no such case in with suture technique. Foroutan et al.<sup>30</sup> reported 20% of cases with graft retraction in his study. Suturing is more time consuming, lead to higher postoperative discomfort, higher recurrence, and complications than others, such as prolonged healing, fibrosis, and granuloma formation.<sup>31,32</sup>

In present study no significant difference was observed between mean vertical (KV) and horizontal (KH) corneal meridian or their difference (KV-KH) by two types of surgery (Pterygium excision with conjunctival autograft without suture or with suture) at various duration of study ( $p > 0.05$ ). It mean in comparison between pre-operative and post-operative status there was equal reduction of KV, difference between KV & KH, increase in KH following two types of surgery. Though, till date, literature search reveals no similar study has been done we saw that as such there is no significant difference in keratometry readings and post operative refraction using these two different techniques, but patient discomfort (i.e. measured by pain score) is relatively less by using sutureless technique rather than doing pterygium surgery with suture.

## 8. Conclusion

Pterygium leads to significant high corneal astigmatism, which hampers vision of the patient. Pterygium causes horizontal meridian flattening (KH). We concluded that, pterygium causes simple myopic with the rule type of astigmatism and compound hypermetropic astigmatism type of refractive error. There is drastic changes in keratometry readings pre-operatively and post surgical excision of pterygium tissue. Overall, after pterygium excision surgery most of the patients had no refractive error. After pterygium excision surgery compound astigmatic type of refractive error was nullified while there was no significant change in simple myopia. Post-operatively, spherical error was reduced and astigmatism was reduced to nullified. With the rule astigmatism was most common pre-operatively. Comparing pre-operative and post-operative refractive error after pterygium excision surgery shows that, in grade 1 pterygium simple myopic astigmatic error almost remains same. In Grade 2, 3 pterygium patients had no refractive error to mild simple myopic astigmatic error but compound hypermetropic astigmatism is completely nullified so patient with grade  $> 2$  pterygium should undergo pterygium excision surgery. We concluded, that with increase in the size of pterygium amount of cornea induced astigmatism also increases. There is definite change in refractive status in pterygium post pterygium surgery so with this study I would like to deduce that, to get emmetropic status for cataract surgery or refractive cataract surgery.

## 9. Source of funding

None.

## 10. Conflict of interest

None.

## References

1. Kanski JJ, Bowling B. Clinical Ophthalmology: A Systematic approach. 7th ed. Elsevier ; 2011,.
2. Lindsay RG, Sullivan L. -. *Clin Exp Optom*. 2001;84(4):200–203.
3. Youngson RM. Pterygium in Israel. *Am J Ophthalmol*. 1972;1:954–959.
4. Varssano D. Etiology and Clinical Presentation of Astigmatism. In *Astigmatism-Optics, Physiology and Management* 2012 Feb 29. Intech Open,.
5. Pavilack MA, Halpern BL. Corneal topographic changes induced by pterygia. *J Refractive Surg*. 1995;11(2):92–95.
6. Buratto L. Corneal topography: The clinical atlas. Slack incorporated; 1996 ;.
7. Lawan SHA, Ifeanyichukwu EP, Yahaya HB, Sani RY, Habib SG, et al. The astigmatic effect of pterygium in a Tertiary Hospital in Kano. *Ann Afr Med*. 2018;17(1):7.
8. Fong KS, Balakrishnan V, Chee SP, Tan DT. Refractive change following pterygium surgery. The CLAO journal: official publication of the Contact Lens Association of Ophthalmologists, Inc. 1998;24(2):115–117.
9. Allan BD, Short P, Crawford GJ, Barrett GD, Constable IJ. Pterygium excision with conjunctival autografting: an effective and safe technique. *Br J Ophthalmol*. 1993;77(11):698–701.
10. Lewallen S. A randomized trial of conjunctival autografting for pterygium in the tropics. *Ophthalmol*. 1989;96(11):1612–1614.
11. Wu KL, He MG, Xu JJ, Li SZ. The epidemiological characteristic of pterygium in middle-aged and the elderly in Doumen County. *J Clin Ophthalmol*. 1999;7(1):17–18.
12. Lemp A. Report of the National Eye Institute/Industry workshop on clinical trials in dry eyes. *Eye Contact Lens*. 1995;21(4):221–232.
13. Pterygium management based upon a theory of pathogenesis. *Trans Am Acad Ophthalmol Otolaryngol*. 1975;79:603–612.
14. Coster D. Pterygium—an ophthalmic enigma. *Br J Ophthalmol*. 1995;79(4):304.
15. Fernandes M, Sangwan VS, Bansal AK, Gangopadhyay N, Sridhar MS, et al. Outcome of pterygium surgery: Analysis over 14 years. *Eye (Lond)*. 2005;19:1182–1190.
16. Kawasaki S, Uno T, Shimamura I, Ohashi Y. Outcome of surgery for recurrent pterygium using intra-operative application of mitomycin C and amniotic membrane transplantation. *Jpn J Ophthalmol*. 2003;47:625–626.
17. Mastropasqua L, Carpineto P, Ciancaglini M, Gallenga PE. Long term results of intraoperative mitomycin C in the treatment of recurrent pterygium. *B Ophthalmol*. 1996;80:288–291.
18. Luanratanakorn P, Ratanapakorn T, Suwan-Apichon O, Chuck RS. Randomised controlled study of conjunctival autograft versus amniotic membrane graft in pterygium excision. *Br J Ophthalmol*. 2006;90(12):1476–1480.
19. Bhargava P, Kochar A, Khan NA, Chandak A, Kumawat S, Garhwal J. Comparison of pre-operative and post-operative astigmatism and visual acuity after pterygium excision followed by sutureless and gluefree conjunctival autograft. *Int J Biomed Res*. 2015;6(10):800–804.
20. Shelke E, Kawalkar U, Wankar R, Nandedkar V, Khaire B, Gosavi V. Effect of pterygium excision on pterygium induced astigmatism and visual acuity. *Age (years)*. 2014;40(9):24–28.
21. Avisar R, Loya N, Yassur Y, Weinberger D. Pterygium-induced corneal astigmatism. *The Israel Med Assoc J: IMAJ*. 2000;2(1):14–15.
22. Lin A, Stern G. Correlation between pterygium size and induced corneal astigmatism. *Cornea*. 1998;17(1):28–30.
23. Popat KB. Keratometry readings before and after pterygium excision surgery. *J Res Med Dent Sci*. 2014;2(3).
24. Kampitak K. The effect of pterygium on corneal astigmatism. *J Med Assoc Thailand= Chotmaihet Thangphaet*. 2003;86(1):16–23.
25. Mohammad-Salih PA, Sharif AF. Analysis of pterygium size and induced corneal astigmatism. *Cornea*. 2008;27(4):434–438.
26. Ali RT, Ali AB. The effect of pterygium on corneal thickness, corneal curvature, tear volume, and intraocular pressure in a Sudanese

- Population. *Al-Basar Int J Ophthalmol*. 2017;4(4):109–109.
27. Maheshwari S. Pterygium-induced corneal refractive changes. *Indian J Ophthalmol*. 2007;55(5):383.
  28. Tomidokoro A, Miyata K, Sakaguchi Y, Samejima T, Tokunaga T, Oshika T. Effects of pterygium on corneal spherical power and astigmatism. *Ophthalmol*. 2000;107(8):1568–1571.
  29. Show J. Clinical Update: Cornea New Approach Emerges for Pterygium Surgery ;. *Eyenet*.
  30. Foroutan A, Beigzadeh F, Ghaempanah MJ, Eshghi P, Amirizadeh N, et al. Efficacy of autologous fibrin glue for primary pterygium surgery with conjunctival autograft. *Iranian J Ophthalmol*. 2011;23(1):39–47.
  31. Hall RC, Logan AJ, Wells AP. Comparison of fibrin glue with sutures for pterygium excision surgery with conjunctival autografts. *Clin Exp Ophthalmol*. 2009;37(6):584–589.
  32. Koranyi G, Seregard S, Kopp ED. Cut and paste: a no suture, small incision approach to pterygium surgery. *Br J Ophthalmol*. 2004;88(7):911–914.

### Author biography

**Bansari Vadodaria** Resident

**Aishwariya Thakre** Assistant Professor

**Rupali Maheshgauri** Professor

**Divya Motwani** Resident

**Ashish Mishra** Resident

**Cite this article:** Vadodaria B, Thakre A, Maheshgauri R, Motwani D, Mishra A. Changes in keratometry and refractive status pre and post pterygium surgery. *Int J Ocul Oncol Oculoplasty* 2019;5(4):205-216.