

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Ocular Oncology and Oculoplasty

Journal homepage: <https://ijooo.org/>

## Editorial

# Oxidative stress and antioxidants in ocular disorders

Rajendra P Maurya<sup>1\*</sup>, Kokil Tiwari<sup>1</sup>

<sup>1</sup>Regional Institute of Ophthalmology, IMS, Banaras Hindu University, Varanasi, Uttar Pradesh, India



## ARTICLE INFO

### Article history:

Received 10-04-2024

Accepted 27-04-2024

Available online 03-05-2024

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

Free radicals are generated during physiological oxidation-reduction reactions being carried out in the powerhouse of cell-mitochondria and endoplasmic reticulum. These free radicals contain unpaired electrons which together with electrophilic molecules are known as reactive oxygen species (ROS) and reactive nitrogen species (RNS).<sup>1,2</sup> Endogenous sources of ROS

are produced as byproducts of various metabolic processes and enzymes: Xanthine oxidase (XO) generates superoxide and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) when oxidizing xanthine to uric acid.<sup>3</sup> Endothelial nitric oxide synthase (eNOS), when uncoupled during endothelial dysfunction, generates superoxide instead of nitric oxide.

Cyclooxygenases (COX), Phospholipase A<sub>2</sub>, Lipoxygenase, and Cytochrome P450 also contribute to ROS production. ROS are also closely linked to inflammation, where immune cells generate excessive ROS to combat pathogens, potentially exacerbating inflammation and tissue damage.<sup>4-6</sup> Exogenous Sources of ROS are air pollution, tobacco smoke and ultraviolet (UV) radiation etc.<sup>7-10</sup>

To counteract the harmful effects of ROS body has antioxidant defense system in place which consists of enzymes like superoxide dismutase (SOD), catalase (CAT), heme oxygenase (HO) and glutathione peroxidase (GPX). Some non-enzymatic scavengers are vitamin E, vitamin C, and glutathione (GSH).<sup>11</sup> Imbalance between generation of

these free radicals and their neutralization by antioxidants leads to oxidative stress. Thus causing damage to various components of the cell by ROS. It causes modification of proteins, lipids and DNA. ROS react with DNA molecules causing oxidation of bases and deoxyribose which causes base modification, rearrangement of DNA sequences, miscoding of DNA, gene duplications or activation of oncogenes. These processes are responsible for carcinogenesis and various other pathologies.<sup>12</sup>

Ocular system with high metabolic activity and permanent exposure to light is extremely susceptible to oxidative stress. It is proposed to be involved in etiopathogenesis of various ocular pathologies like dry eye, pterygium, cataract, keratoconus, Fuchs endothelial corneal dystrophy, diabetic retinopathy, age related macular degeneration, retinal vein occlusion, retinopathy of prematurity and ocular malignancies etc.<sup>13,14</sup>

## 1. Dry Eye Diseases

It is a multifactorial disorder. Oxidative stress has a major contribution to its development and progression. A meta-analysis revealed that patients with Dry Eye Disease (DED) tend to have higher levels of oxidative stress markers in their tear fluid and conjunctival specimens, indicating a significant link between oxidative stress and DED. The oxidative stress markers that were found to be elevated include lipid peroxide, myeloperoxidase, nitric oxide (NO) synthase, xanthine oxidase/oxidoreductase, 4-HNE, and

\* Corresponding author.

E-mail address: [mauryarp\\_bhu@yahoo.com](mailto:mauryarp_bhu@yahoo.com) (R. P. Maurya).

malondialdehyde (MDA). These markers suggest increased lipid oxidation, ROS production, and oxidative damage in DED.<sup>15</sup>

Furthermore, the study noted that patients with DED exhibit impaired antioxidant functions. In particular, the expression of key antioxidant enzymes such as catalase (CAT), glutathione peroxidase (GPX), and superoxide dismutase (SOD) was reduced in conjunctival epithelial cells compared to healthy controls.<sup>16</sup> This reduced expression of antioxidant enzymes suggests that DED may involve not only an increase in oxidative stress but also a decreased capacity to neutralize ROS, potentially contributing to the disease's progression and severity. Aging is one of the risk factors for DEDs which causes an increase in oxidative stress due to declining antioxidant defense mechanism.

ROS induce inflammatory cytokines such as IL-1, IL-6 and TNF-alpha which are responsible for DED. These levels correlate with disease severity.<sup>17-19</sup>

Decreased levels of SOD expression with elevated oxidative stress markers were noted in mice with meibomian gland dysfunction which is major factor behind tear film instability.<sup>20</sup>

Oxidative damage to lacrimal gland has also been proposed for the etiopathogenesis of DED.

Therapeutic approach to manage oxidative stress in DED is by giving antioxidants like vitamin A and E, and coenzyme Q10 topically in the form of tear substitutes.<sup>21</sup> Lactoferrin was also found to have a beneficial role in treatment of DED.<sup>22</sup> Iodide iontophoresis uses electrical current to drive iodide ions into body tissues. This technique has been applied to the eye to increase the antioxidant capacity of tear fluid, potentially benefiting patients with DED. Studies suggest that it can improve symptoms, tear film stability, and reduce morphological changes in DED patients. However, its use is limited by the potential cytotoxic effects of iodide on corneal epithelial and endothelial cells.<sup>23-25</sup>

## **2. Pterygium**

Oxidative stress due to phototoxic effects of UV radiation plays a crucial role in pathogenesis of pterygium. Significant higher levels of oxidative DNA damage marker 8-OHdG were found in pterygium specimens.<sup>26</sup>

Antioxidants have been identified as a potential therapeutic tool for treating pterygium. Phenolic compounds, such as flavonoids, carotenoids, curcumin, ellagic acid, and chalcones, are known for their anti-inflammatory and antioxidant properties.<sup>27</sup> Curcumin, in particular, has shown promising results in reducing the proliferation of pterygium fibroblasts.<sup>28</sup> Other therapeutic strategies focus on targeting ROS-related pathways and signaling molecules that contribute to pterygium. For instance, bevacizumab, a recombinant humanized anti-

VEGF antibody, is used in perioperative subconjunctival injections to lower the recurrence rate of pterygium.<sup>29-31</sup> Additionally, cyclosporine A, an immunosuppressive agent, has demonstrated the ability to suppress cell migration and reduce the expression of MMP-3 and MMP-13, resulting in fewer recurrences when used as an adjuvant with pterygium excision.<sup>32</sup>

## **3. Keratoconus**

In a meta-analysis by Navel et al, a marked decrease in antioxidant levels was noted in keratoconus patients. Increased levels of proinflammatory cytokines like IL-6, TNF-alpha and MMP were seen. This imbalance leads to keratocyte apoptosis and changes in the corneal extra cellular matrix leading to disease progression.<sup>33</sup> Flavonoids have shown to protect against UV rays induced stress. Lactoferrin loaded contact lenses are also one of the therapeutic options available. Vitamin D supplementation helps in reduction of inflammation and oxidative stress.<sup>34,35</sup>

## **4. Cataract**

The ongoing metabolic processes in lens lead to formation of reactive oxygen species (ROS), which may harm lens cells and increase the risk of cataracts. Additionally, sunlight exposure can lead to ROS formation through photoexcitation, where absorbed light shifts from a photoexcited singlet state to an excited triplet state, causing ROS generation.<sup>36</sup>

To counteract these harmful effects, the lens employs several antioxidants and protective mechanisms. Glutathione peroxidase and catalase neutralize hydrogen peroxide and lipid peroxide, reducing oxidative stress. Superoxide dismutase (SOD) converts superoxide radicals into less harmful substances, and ascorbate (vitamin C) in the lens's outer layers neutralizes superoxide, peroxide, and hydroxyl ions while also scavenging singlet oxygen and thiol radicals.

Additionally, the lens has natural protection against ultraviolet (UV) radiation through aromatic amino acids like tryptophan, fluorophores, and 3-hydroxykynurenine, which absorb and filter UV rays, preventing UV-induced damage.<sup>37</sup> These antioxidants and protective mechanisms help maintain the clarity and health of the lens by reducing oxidative stress and minimizing the effects of ROS. Supplementation of vitamin E, lutein, zeaxanthin or carotenoids is related to reduced risk for age-related cataract.

## **5. Age Related Macular Degeneration**

Excessive production of reactive oxygen species (ROS) can activate cellular signaling pathways that drive inflammation and other harmful effects. ROS can stimulate the mitogen-activated protein kinases (MAPKs) and nuclear factor

kappa B (NF- $\kappa$ B) pathways, leading to higher levels of inflammatory cytokines like interleukin-1 beta (IL-1 $\beta$ ), interleukin-18 (IL-18), and tumor necrosis factor alpha (TNF- $\alpha$ ). This is mediated through the NLRP3 inflammasome and caspase-1<sup>38</sup>. ROS also influence angiogenesis by increasing vascular endothelial growth factor (VEGF) via hypoxia-inducible factor alpha (HIF- $\alpha$ ). Higher VEGF levels can cause abnormal blood vessel growth and choroidal vascular dysfunction. Additionally, ROS can reduce nitric oxide (NO) levels, diminishing its vasoprotective role, which further contributes to vascular problems.<sup>39</sup>

Anti VEGF therapy has shown to have major role in its treatment. Lutein and zeaxanthin are well-known carotenoids that support eye health, particularly in reducing the risk of age-related macular degeneration (AMD). The Age-Related Eye Disease Study (AREDS) showed that daily supplementation could help slow AMD progression. The supplement regimen consisted of a combination of vitamin C, vitamin E, cupric oxide, zinc, and beta-carotene, which are antioxidants that protect against oxidative stress. This combination was found to be effective in reducing the risk of AMD progression in the study's participants. Novel treatment targeting P62 and Nrf2 have shown beneficial effects.<sup>40,41</sup>

## **6. Ocular Malignancies**

Cancer is a complex process with three main stages: initiation, promotion, and progression. Oxidative stress, caused by an imbalance of reactive oxygen species (ROS) and antioxidants, plays a role in all three stages. In the initiation stage, ROS can damage DNA, leading to gene mutations and structural changes that initiate cancer development. During the promotion stage, ROS may disrupt gene expression, cell-to-cell communication, and second messenger systems, leading to increased cell proliferation and decreased apoptosis, encouraging the growth of initiated cells. In the progression stage, oxidative stress can cause further DNA alterations and genomic instability, promoting more aggressive tumor characteristics and the potential for metastasis.

This interaction with oxidative stress across all stages underscores its role in cancer development and progression, indicating the importance of maintaining a healthy balance between ROS and antioxidants to help mitigate cancer risk.<sup>42,43</sup>

Antioxidants have gained significant interest in cancer therapy due to their ability to neutralize oxidizing free radicals, which helps prevent cellular damage during chemotherapy. The use of antioxidants in chemotherapy aims to protect normal tissues from oxidative damage without negatively impacting the effectiveness of tumor-targeting treatments. Research on antioxidant supplementation during chemotherapy has shown that

there are two approaches to using antioxidants in cancer therapy. (i) Preventive low doses protect both normal and tumor cells from oxidative damage, potentially reducing side effects without impacting chemotherapy outcomes. (ii) Therapeutic high doses of antioxidants can inhibit the growth of cancer cells while sparing normal cells, contributing to enhanced cancer treatment. Recent reviews have highlighted the following benefits of administering antioxidants alongside chemotherapy:

Antioxidants at specific doses do not disrupt the efficacy of chemotherapy in targeting cancer cells. Improved cytotoxic effects: Antioxidants can enhance the destructive effects of chemotherapy on cancer cells while providing a protective barrier for normal tissues. This can lead to increased patient survival and better therapeutic responses.<sup>44</sup>

Overall, the integration of antioxidants into cancer therapy shows promise for reducing chemotherapy-induced damage to normal tissues and potentially improving cancer treatment outcomes. However, further research is needed to understand the optimal types and doses of antioxidants for safe and effective use in cancer therapy.

Various studies have investigated potential therapeutic agents to combat oxidative stress in ocular diseases, with a focus on antioxidants and other related compounds. Antioxidant supplementation, such as with vitamins A and E, has been examined in numerous studies, but the results have been contradictory. This inconsistency suggests that while antioxidants can play a role in reducing oxidative stress, their effectiveness may depend on specific conditions or individual responses.

One promising agent in this field is idebenone, a synthetic analogue of Coenzyme Q<sub>10</sub> (CoQ<sub>10</sub>). Approved by the European Medicine Agency (EMA) in 2015, idebenone is the only disease-specific medication for Leber's Hereditary Optic Neuropathy (LHON), a genetic condition leading to vision loss. Its approval represents a significant advancement in targeted treatments for specific ocular diseases related to oxidative stress.

Another key area of interest involves the Nrf2 (Nuclear factor erythroid 2-related factor 2) pathway. Nrf2 is a crucial transcription factor that regulates antioxidant enzyme expression, playing an essential role in cellular defense against oxidative stress. Disturbances in Nrf2 function have been implicated in several ocular diseases, making the modulation of this pathway a promising therapeutic approach. By enhancing Nrf2 activity, it may be possible to increase the body's natural antioxidant defense mechanisms, thereby reducing oxidative stress-related damage in the eye.<sup>45</sup>

Despite these promising leads, it's important to note that while experimental studies have yielded encouraging results, comprehensive clinical trials are necessary to confirm the safety and effectiveness of these therapeutic

approaches. These studies are crucial to determining the appropriate dosages, long-term effects, and specific applications for different ocular diseases influenced by oxidative stress.

## 7. Conflict of Interest


None.

## References

- Wang J, Li M, Geng Z, Khattak S, Ji X, Wu D, et al. Role of oxidative stress in retinal disease and the early intervention strategies: a review. *Oxid Med Cell Longev*. 2022;doi:10.1155/2022/7836828.
- Lushchak VI. Classification of oxidative stress based on its intensity. *Excli J*. 2014;13:922–37.
- Förstermann UF, Xia N, Li H. Roles of Vascular Oxidative Stress and Nitric Oxide in the Pathogenesis of Atherosclerosis. *Circ Res*. 2017;120(4):713–35.
- Coon MJ, Vaz AD, Mcginnity DF, Peng HM. Multiple activated oxygen species in P450 catalysis: contributions to specificity in drug metabolism. *Drug Metab Dispos*. 1998;26(12):1190–3.
- Wang Y, Zhang SXL, Gozal D. Reactive Oxygen Species and the Brain in Sleep Apnea. *Respir Physiol Neurobiol*. 2010;174(3):307–16.
- Dammak A, Pastrana C, Martin-Gil A, Carpena-Torres C, Cerda A, Simovart M, et al. Oxidative stress in the anterior ocular diseases: diagnostic and treatment. *Biomedicines*. 2023;11(2):292. doi:10.3390/biomedicines11020292.
- Kuntic M, Kuntic I, Krishnankutty R, Gericke A, Oelze M, Junglas T, et al. Co-exposure to urban particulate matter and aircraft noise adversely impacts the cerebro-pulmonary-cardiovascular axis in mice. *Redox Biol*. 2023;59:102580. doi:10.1016/j.redox.2022.102580.
- Hahad O, Kuntic M, Kuntic I, Daiber A, Münzel T. Tobacco smoking and vascular biology and function: evidence from human studies. *Pflügers Archiv*. 2023;475(7):797–805.
- Guo C, Ning X, Zhang J, Zhang C, Wang J, Su L, et al. Ultraviolet B radiation induces oxidative stress and apoptosis in human lens epithelium cells by activating NF- $\kappa$ B signaling to down-regulate sodium vitamin C transporter 2 (SVCT2) expression. *Cell Cycle*. 2023;22(12):1450–62.
- Jimenez MTB, Gericke A, Frenis K, Rajlic S, Kvandova M, Kröllerschön S, et al. Effects of aircraft noise cessation on blood pressure, cardio- and cerebrovascular endothelial function, oxidative stress, and inflammation in an experimental animal model. *Sci Total Environ*. 2023;903:166106. doi:10.1016/j.scitotenv.2023.166106.
- Brieger K, Schiavone S, Müller FJ, Krause KH. Reactive oxygen species: from health to disease. *Swiss Med Wkly*. 2012;142:13659. doi:10.4414/sm.w.2012.13659.
- Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran pathologic basis of disease. Philadelphia: Elsevier health sciences (p)Ltd.; p. 52–4.
- Hsueh YJ, Chen YN, Tsao YT, Cheng CM, Wu WC, Chen HC, et al. The pathomechanism, antioxidant biomarkers, and treatment of oxidative stress-related eye diseases. *Int J Mol Sci*. 2022;23(3):1255. doi:10.3390/ijms23031255.
- Vallabh NA, Romano V, Willoughby CE. Mitochondrial dysfunction and oxidative stress in corneal disease. *Mitochondrion*. 2017;36:103–13. doi:10.1016/j.mito.2017.05.009.
- Navel V, Sapin V, Henrioux F, Blanchon L, e AL, Chiambaretta F. Oxidative and antioxidative stress markers in dry eye disease: a systematic review and meta-analysis. *Acta Ophthalmol*. 2022;100(1):45–57.
- Cejková J, Ardan T, Simonová Z, Cejka C, Malec J, Dotrelová D, et al. Decreased expression of antioxidant enzymes in the conjunctival epithelium of dry eye (Sjögren's syndrome) and its possible contribution to the development of ocular surface oxidative injuries. *Histol Histopathol*. 2008;23(12):1477–83.
- Enríquez-De-Salamanca A, Castellanos E, Stern ME, andez IF, Carreño E, García-Vázquez C, et al. Tear cytokine and chemokine analysis and clinical correlations in evaporative-type dry eye disease. *Mol Vis*. 2010;16:862–73.
- Yoon KC, Jeong IY, Park YG, Yang SY. Interleukin-6 and tumor necrosis factor-alpha levels in tears of patients with dry eye syndrome. *Cornea*. 2007;26(4):431–7.
- Maurya RP, Gupta S, Ranjan S. Biomarkers of oxidative stress and inflammation in dry eye diseases. *IP Int J Ocul Oncol Oculoplasty*. 2023;9(4):155–6.
- Ibrahim OM, Dogru M, Matsumoto Y, Igarashi A, Kojima T, Wakamatsu TH, et al. Oxidative stress induced age dependent meibomian gland dysfunction in Cu, Zn-superoxide dismutase-1 (Sod1) knockout mice. *PLoS One*. 2014;9(7):e9932. doi:10.1371/journal.pone.0099328.
- Labetoulle M, Benitez-Del-Castillo JM, Barabino S, Vanrell R, Daull P, Garrigue JS, et al. Artificial tears: biological role of their ingredients in the management of dry eye disease. *Int J Mol Sci*. 2022;23(5):2434. doi:10.3390/ijms23052434.
- Versura P, Bavelloni A, Grillini M, Fresina M, Campos EC. Diagnostic performance of a tear protein panel in early dry eye. *Mol Vis*. 2013;19:1247–57.
- Seen S, Tong L. Dry eye disease and oxidative stress. *Acta Ophthalmol*. 2018;96(4):412–20.
- Rieger G, Klieber M, Schimetta W, olz WP, Griebenow S, Winkler R, et al. The effect of iodide iontophoresis on the antioxidative capacity of the tear fluid. *Graefes Arch Clin Exp Ophthalmol*. 2010;248(11):1639–46.
- Shibata Y, Tanaka Y, Tomita T, Taogoshi T, Kimura Y, Chikama T, et al. Evaluation of corneal damage caused by iodine preparations using human corneal epithelial cells. *Jpn J Ophthalmol*. 2014;58(6):522–7.
- Kilic-Toprak E, Toprak I, Caliskan S, Ozdemir Y, Demirtas O, Altintas F, et al. Oxidative stress and genotoxicity in pterygium: a systemic investigation. *Eye Contact Lens*. 2019;45(6):399–404.
- Baهران SS, Alany RG, Schwikkard S, Muen W, Salman LN, Freestone N, et al. Pharmacological treatment strategies of pterygium: Drugs, biologics, and novel natural products. *Drug Discov Today*. 2023;28(1):103416. doi:10.1016/j.drudis.2022.103416.
- Lu CW, Hao JL, Yao L, Li HJ, Zhou DD. Efficacy of curcumin in inducing apoptosis and inhibiting the expression of VEGF in human pterygium fibroblasts. *Int J Mol Med*. 2017;39(5):1149–54.
- neda ANC, Olvera-Morales O, Ramos-Castellon C, Garnica-Hayashi L, Garfias Y. Randomized, controlled trial of conjunctival autografting combined with subconjunctival bevacizumab for primary pterygium treatment: 1-year follow-up. *Clin Exp Ophthalmol*. 2014;42(3):235–41.
- Yang HK, Lee YJ, Hyon JY, Kim KG, Han SB. Efficacy of bevacizumab injection after pterygium excision and limbal conjunctival autograft with limbal fixation suture. *Graefes Arch Clin Exp Ophthalmol*. 2020;258(7):1451–7.
- Sayadi J, Gouider D, Henchiri M, Choura R, Boujelbene N, Abbes I, et al. Preoperative intralesional bevacizumab injection in primary pterygium in Tunisian patients: a randomized controlled prospective study. *J Curr Ophthalmol*. 2022;34(3):297–304.
- Kim YH, Jung JC, Jung SY, Kim YI, Lee KW, Park YJ, et al. Cyclosporine A downregulates MMP-3 and MMP-13 expression in cultured pterygium fibroblasts. *Cornea*. 2015;34(9):297–304.
- Navel V, Malecaze J, Pereira B, Baker JS, Malecaze F, Sapin V, et al. Oxidative and antioxidative stress markers in keratoconus: a systematic review and meta-analysis. *Acta Ophthalmol*. 2021;99(6):777–94.
- Mckay TB, Karamichos D. Quercetin and the ocular surface: What we know and where we are going. *Exp Biol Med*. 2017;242(6):565–72.
- Vitar RM, Fonteyne P, Knutsson KA, Bertuzzi F, Galli L, Rama P, et al. Vitamin D supplementation impacts systemic biomarkers of collagen degradation and copper metabolism in patients with keratoconus. *Transl Vis Sci Technol*. 2022;11(12):16. doi:10.1167/tvst.11.12.16.
- Khurana AK. Disorders of lens and cataract surgery. 1 edn. New Delhi: CBS publishers (p) Ltd.; p. 59–65.

37. Yanoff M, Duckers J. Ophthalmology. 4 edn.. vol. 4. China: Elsevier Saunders publishers (p) Ltd; 2014. p. 330.e1–e11.
38. Ruan Y, Jiang S, Gericke A. Age-Related Macular Degeneration: Role of Oxidative Stress and Blood Vessels. *Int J Mol Sci.* 2021;22(3):1296. doi:10.3390/ijms22031296.
39. Monaghan-Benson E, Hartmann J, Vendrov AE, Budd S, Byfield G, Parker A, et al. The role of vascular endothelial growth factor-induced activation of NADPH oxidase in choroidal endothelial cells and choroidal neovascularization. *Am J Pathol.* 2010;177(4):2091–102.
40. Li LH, Lee JC, Leung HH, Lam WC, Fu Z, Lo CY, et al. Lutein Supplementation for Eye Diseases. *Nutrients.* 2020;12(6):1721. doi:10.3390/nu12061721.
41. Bellezza I. Oxidative stress in age-related macular degeneration: Nrf2 as therapeutic target. *Front Pharmacol.* 2018;9:1280. doi:10.3389/fphar.2018.01280.
42. Klaunig JE, Xu Y, Isenberg JS, Bachowski S, Kolaja KL, Jiang J, et al. Jr The role of oxidative stress in chemical carcinogenesis. *Environ Health Perspect.* 1998;106(1):289–5.
43. Maurya RP, Prajapat MK, Singh VP, Roy M, Todi R, Bosak S, et al. Serum Malondehyde as a Biomarker of oxidative stress in patient with primary ocular carcinoma: Impact on response to chemotherapy. *Clin Ophthalmol.* 2021;p. 871–9. doi:10.2147/OPHTH.S287747.
44. Ammar HO, Shamma RN, Elbatanony RSE, Khater B, Ammar HO, Shamma RN, et al. Antioxidants in Cancer Therapy: Recent Trends in Application of Nanotechnology for Enhanced Delivery. *Sci Pharm.* 2020;88(1). doi:10.3390/scipharm88010005.
45. Böhm EW, Buonfiglio F, Voigt AM, Bachmann P, Safi T, Pfeiffer N, et al. Oxidative stress in the eye and its role in the pathophysiology of ocular diseases. *Redox Biol.* 2023;68:102967. doi:10.1016/j.redox.2023.102967.

### Author biography

**Rajendra P Maurya**, Associate Professor  <https://orcid.org/0000-0001-9343-6003>

**Kokil Tiwari**, Junior Resident

**Cite this article:** Maurya RP, Tiwari K. Oxidative stress and antioxidants in ocular disorders. *IP Int J Ocul Oncol Oculoplasty* 2024;10(1):1-5.